

POLICY: 554.62
TITLE: Hypothermia - Frostbite

EFFECTIVE: 07/01/2024
REVIEW: 07/2027
SUPERCEDES:

APPROVAL SIGNATURES ON FILE IN EMS OFFICE

PAGE: 1 of 2

HYPOTHERMIA - FROSTBITE

- I. AUTHORITY
Health and Safety Code, Division 2.5, California Code of Regulations, Title 22, Division 9
- II. PURPOSE
To serve as a patient treatment standard for EMRs, EMTs, and Paramedics within their scope of practice.
- III. PROTOCOL
Patients with severe hypothermia may appear dead (pulseless, absent respiration, and fixed pupils) but still have cardiac electrical activity.

Provider Key: F = First Responder/EMR E = EMT O = EMT Local Optional SOP
P = Paramedic D = Base Hospital Physician Order Required

USE EXTREME CAUTION WHEN MOVING PATIENT

Moderate Hypothermia (92°-95° F / 33°-35° C) Severe Hypothermia (Core temp < 92° F / < 33° C)	F	E	O	P	D
ASSESSMENT	X	X	X	X	
PULSE OXIMETRY: apply and monitor.		X	X	X	
CAPNOGRAPHY: apply and monitor.				X	
ADVANCED AIRWAY: if GCS is < 8 and not rapidly improving, consider: - SGA - or ETI			X	X X	
OXYGEN: if pulse oximetry <94% or signs of respiratory distress or hypoperfusion.	X	X	X	X	
WARMING MEASURES: remove wet clothing and cover with warm dry blankets. Use ambient heat and heat packs as able.	X	X	X	X	
ECG MONITOR: lead placement may be delegated. Treat as indicated.				X	
VASCULAR ACCESS: IV/IO, rate as indicated with warm fluids.				X	
TEST FOR GLUCOSE		X	X	X	
ORAL GLUCOSE: consider administering oral glucose to patients who are awake and have an intact gag reflex.		X	X	X	
D10: infuse 100 mL IV/IO if blood glucose < 70 mg/dL. Recheck blood glucose 10 minutes post infusion. If blood glucose < 70 mg/dL infuse remaining 150 mL.				X	
GLUCAGON: If no IV/IO access and unable to tolerate oral glucose, give 1 mg IM if blood glucose < 70 mg/dL. Recheck blood glucose 10 minutes post injection. If blood glucose remains < 70 mg/dL, repeat 1 mg IM.				X	

FROSTBITE					
(skin is white, numb or burning, soft to touch and does not re-color with touch)					
	F	E	O	P	D
EVALUATE: for hypothermia and treat as indicated.	X	X	X	X	
WARMING MEASURES: move patient to warm environment & wrap affected extremity with thick, warmed blankets or clothing. DO NOT RUB AFFECTED EXTREMITY & AVOID CHEMICAL HEAT PACKS.	X	X	X	X	
PAIN MANAGEMENT: Refer to 554.44 PAIN MANAGEMENT as indicated.				X	