

POLICY: 555.52
TITLE: Pediatric Dystonic Reactions to Phenothiazines

EFFECTIVE: 02/01/2026DRAFT
REVIEW: 02/2028DRAFT
SUPERCEDES:

APPROVAL SIGNATURES ON FILE IN EMS OFFICE

PAGE: 1 of 1

PEDIATRIC DYSTONIC REACTIONS TO PHENOTHIAZINES

- I. AUTHORITY
Health and Safety Code, Division 2.5, California Code of Regulations, Title 22, Division 9
- II. PURPOSE
To serve as a patient treatment standard for EMRs, EMTs, AEMT's, and Paramedics within their scope of practice.
- III. PROTOCOL

Provider Key: F = First Responder/EMR E = EMT O = EMT Local Optional SOP
P = Paramedic A=Advanced EMT ——— D = Base Hospital Physician Order
Required

	F	E	O	<u>A</u>	P	D
ASSESSMENT	X	X	X	<u>X</u>	X	
BLS AIRWAY: okay if airway patent. Support ventilations with appropriate airway adjuncts.	X	X	X	<u>X</u>	X	
SUPRAGLOTTIC AIRWAY: if GCS is < 8 and not rapidly improving, consider SGA.			X	<u>X</u>	X	
PULSE OXIMETRY: apply and monitor.		X	X	<u>X</u>	X	
CAPNOGRAPHY: apply and monitor if SGA has been placed.					X	
OXYGEN: if pulse oximetry < 94% or signs of respiratory distress or hypoperfusion.	X	X	X	<u>X</u>	X	
ECG MONITOR: lead placement may be delegated. Treat as indicated.					X	
VASCULAR ACCESS: IV/IO, rate as indicated.				<u>X</u>	X	
DIPHENHYDRAMINE: 1 mg/kg IV/IO push titrated to relief of signs and symptoms or IM if IV/IO access not promptly available. Max dose 50 mg.					X	
ACTIVATED CHARCOAL: 1 g/kg, maximum 50 g PO if patient's GCS is 15 or via NG if patient is intubated and oral ingestion has occurred with 60 minutes. (FOR OVERDOSE VIA INGESTION ONLY). <u>AEMT PO administration is PO only.</u>				<u>X</u>	X	
NASOGASTRIC TUBE: suction gastric contents only if patient has SGA and oral ingestion has occurred with 60 minutes. (FOR OVERDOSE VIA INGESTION ONLY).					X	

NOTE: Phenothiazine reactions may occur at normal dosing levels and the induction of vomiting is not recommended.

Pediatric Normal Vital Signs					
<i>Age</i>	HR	RR	BP	Temp (C)	Temp (F)
<i>Premie</i>	120-170	40-70	55-75/35-45	36-38	96.8-100.4
<i>0-3 months</i>	100-160	35-60	65-85/45-55	36-38	96.8-100.4
<i>3-6 months</i>	90-120	30-45	70-90/50-65	36-38	96.8-100.4
<i>6-12 months</i>	80-120	25-40	80-100/55-65	36-38	96.8-100.4
<i>1-3 years</i>	70-110	20-30	90-105/55-70	36-38	96.8-100.4
<i>3-6 years</i>	65-110	20-25	90-110/60-75	36-38	96.8-100.4
<i>6-12 years</i>	65-100	14-22	90-120/60-75	36-38	96.8-100.4
<i>12+</i>	55-100	12-20	100-135/65-85	36-38	96.8-100.4